

Request for Information Reportable/Communicable Disease – Co-infected Chlamydia and Gonorrhoea

Information on this form is collected under the Health Protection and Promotion Act, R.S.O. 1990.

Date: _____

Clinic/Practitioner: _____

Fax: _____

From: Clinical Services

Tel: 1-800-265-7293 ext. 4556

Fax: 1-855-934-5463

PLEASE COMPLETE FORM AND RETURN AS SOON AS POSSIBLE

Patient Name: _____ D.O.B.: _____
Address: _____
Tel: _____ Cell: _____

Patient demographics correct?	
<input type="checkbox"/> Yes	<input type="checkbox"/> No

Reason for Testing

Routine: Yes No

Contact of Case: Yes No

Symptomatic: Yes No If yes, please list _____

Patient pregnant: Yes No If yes, please provide E.D.D. _____

Patient co-infection: HIV: Yes No VDRL: Yes No

Patient advised of result: Yes No

Patient advised to notify possible partners from the last 60 days: Yes No

Multiple sex partners: Yes No Same sex partners: Yes No

Who will notify partners? Patient (case) Health care provider

Partner notification: All partners 60 days prior to diagnosis advised to seek testing and treatment

First Line Treatment – when co-infected with both Chlamydia and Gonorrhoea

Azithromycin 1 gm AND Ceftriaxone 500 mg IM (Monotherapy) Date: _____

Alternative Treatment – refer to Canadian Guidelines on Sexually Transmitted Infections

Other: _____ Date: _____

Counselling Provided

TEST OF CURE:

Gonorrhoea testing: Recommended for all positive GC cases at all positive sites

- culture 3-21 days after completing treatment.
- Asymptomatic – NAAT >3 weeks after completing treatment.
- Symptomatic – NAAT and Culture preferred
- RETESTING – due to the risk of re-infection, all GC cases should be re-screened 6 months following treatment.

Chlamydia testing: recommended for pregnant, second line treatment, rectal/pharyngeal, suspected non-compliance.

- NAAT – 4-6 weeks after completing treatment

Patient advised to abstain from all sexual activity (oral, anal and vaginal) for 7 days post treatment. Safer sex practices discussed, e.g. use of condoms.

Azithromycin, Doxycycline and Ceftriaxone are available free of charge through Public Health.

To order your free STI medication, please download an STI order form from our website.

IMPORTANT: A PHN will attempt to contact client for counselling, partner notification and future infection prevention. We will also confirm that the client took and tolerated the medication you prescribed. We will discuss test of cure and other possible STI testing if appropriate post treatment.



Public Health
WELLINGTON-DUFFERIN-GUELPH
Stay Well.

1-800-265-7293 | www.wdgppublichealth.ca | info@wdgppublichealth.ca