

ADDITIONAL PRECAUTIONS

IPAC MEASURES TO FOLLOW AND SIGNAGE TO USE

Assess the individual

- Perform hand hygiene
 - Complete a point of care risk assessment (PCRA)
 - Wear appropriate PPE
- Do they have new or worsening RESPIRATORY symptoms?**
- Cough
 - Sore throat
 - Fever
 - Congestion

YES

3

NO

Do they have gastrointestinal symptoms?

- Nausea
- Vomiting
- Diarrhea

YES

2

NO

1

1



ROUTINE PRACTICES

- Remember the following before every interaction:
 - Point of care risk assessment (PCRA)
 - Hand hygiene
 - Wear personal protective equipment (PPE) based on your PCRA
- Assess the individual for any respiratory or gastrointestinal symptoms

2



CONTACT

- STAFF: gown and gloves
- Hand hygiene- soap and water is preferred if handling loose stool
- Dedicate medical equipment and/or clean and disinfect between use



3



DROPLET CONTACT

- STAFF: medical mask, eye protection, gown and gloves
- If COVID-19 is suspected or confirmed, either an N95 respirator or well fitting medical mask is recommended
- Dedicate medical equipment and/or clean and disinfect between use
- INDIVIDUAL: medical mask if outside the room (if tolerated)

