

Recommended Immunization Schedule for Children from Infancy to 6 Years



Given by
needle



Given by
oral drops

Age	Vaccines	Abbreviation
2 months	 Diphtheria, tetanus, pertussis, polio, haemophilus influenza type b vaccine	DTaP-IPV-Hib
	 Pneumococcal conjugate vaccine	Pneu-C-13
	 Rotavirus vaccine	Rot-1
4 months	 Diphtheria, tetanus, pertussis, polio, haemophilus influenza type b vaccine	DTaP-IPV-Hib
	 Pneumococcal conjugate vaccine	Pneu-C-13
	 Rotavirus vaccine	Rot-1
6 months	 Diphtheria, tetanus, pertussis, polio, haemophilus influenza type b vaccine	DTaP-IPV-Hib
12 months	 Measles, mumps, rubella vaccine <i>2 doses are required for school attendance*; must be given on or after the 1st birthday and at least 28 days apart</i>	MMR
	 Meningococcal conjugate vaccine <i>Must be given on or after the 1st birthday</i>	Men-C-C
	 Pneumococcal conjugate vaccine	Pneu-C-13
15 months	 Varicella (chickenpox) vaccine	Var
18 months	 Diphtheria, tetanus, pertussis, polio, haemophilus influenza type b vaccine	DTaP-IPV-Hib
4-6 years	 Tetanus, diphtheria, pertussis, polio vaccine	Tdap-IPV
	 Measles, mumps, rubella, varicella (chickenpox) vaccine <i>*The second dose of MMR is usually given in combination with the second recommended dose of varicella at 4-6 years</i>	MMRV