


DILUTE BEFORE USE




- Remove a thawed vial of Pfizer-BioNTech COVID-19 Vaccine from the refrigerator and allow it to come to room temperature.
- If using a frozen vial of Pfizer-BioNTech COVID-19 Vaccine, thaw for 30 minutes at room temperature.


Vials at room temperature must be diluted within 2 hours.

Requires 15 minutes to come to room temperature


Time at room temperature is cumulative. Unmixed vaccine may be returned to the fridge if not needed right away. However, any time at room temperature needs to be marked on the vial. The next time it is brought out of the fridge it then only has the 2 hours minus the original time out before it must be diluted. For example, if vaccine was out of fridge for .5 hour and then returned to fridge; when vaccine is next removed it must be mixed within 1.5 hours.



- Before dilution, invert **gently** 10 times to mix.
- Do not shake.



- Obtain sterile 0.9% Sodium Chloride Injection, USP.
- Cleanse the vial stopper with a single-use antiseptic swab.
- Add 1.8 mL of 0.9% Sodium Chloride Injection, USP into the Pfizer-BioNTech COVID-19 Vaccine vial using a needle 21-gauge or narrower.



- Record the date and time of dilution on the Pfizer-BioNTech COVID-19 Vaccine vial label.
- Store between 2°C to 25°C (35°F to 77°F).
- Discard any unused vaccine 6 hours after dilution.

<p>Pull back plunger to 1.8 mL to remove air from vial</p>	<ul style="list-style-type: none"> • Equalize vial pressure before removing the needle from the vial by withdrawing 1.8 mL air into the empty diluent syringe.
<p>Gently x 10</p>	<ul style="list-style-type: none"> • Gently invert the vial containing the Pfizer-BioNTech COVID-19 Vaccine 10 times to mix. • <u>Do not shake.</u>

<p>PREPARATION OF INDIVIDUAL 0.3 mL DOSES OF PFIZER-BIONTECH COVID-19 VACCINE</p>	
<p>0.3 mL diluted vaccine</p>	<ul style="list-style-type: none"> • Using aseptic technique, cleanse the vial stopper with a single-use antiseptic swab, and withdraw <u>0.3 mL</u> of the Pfizer-BioNTech COVID-19 Vaccine. • Administer immediately, and no later than 6 hours after dilution.

When drawing up the 0.3 mL – keep the needle in the vial to remove any air bubbles or air from the syringe.

DO NOT tap or flick at the syringe to remove bubbles. Handle syringe gently and remember to keep protected from light.